

Light-Cone Gauge Quantization of String Theories on AdS_3 Space

MING YU ¹ and BO ZHANG ²

Institute of Theoretical Physics, Academia Sinica

P.O.Box 2735, Beijing 100080, P.R.China

Abstract

Light-cone gauge quantization procedures are given, for superstring theory on AdS_3 space charged with NS-NS background, both in NSR and GS formalism. The spacetime (super)conformal algebras are constructed in terms of the transversal physical degrees of freedom. The spacetime conformal anomaly agrees with that of covariant formalism, provided that the worldsheet conformal anomaly c equals 26 or 15 for bosonic string or superstring, respectively. The spacetime (super)conformal field theory is found to correspond to orbifold construction on symmetric product space $Sym_p \mathcal{M}/Z_p$.

¹e-mail: yum@itp.ac.cn

²e-mail: zhangb@itp.ac.cn

1 Introduction

Recently, String theories on AdS^3 space has attracted much attention [4, 5, 6, 8, 9, 18, 20, 21, 22, 23, 24]. In ref. [1], Maldacena has conjectured that a type IIB superstring theory on AdS_{d+1} is dual to a superconformal field theory living on the d-dimensional boundary of AdS_{d+1} , which is a compactified Minkowski space. A more precise statement on the AdS/CFT correspondence is give in refs. [2, 3], which relate the boundary values in the former theory to the source terms in the later. Some well studied examples of the AdS/CFT correspondence are the cases of AdS_5 and AdS_3 . On each boundary resp., one finds a 4d $N = 4$ superconformal Yang-Mills theory in the former case, and a 2d $N = 4$ superconformal field theory in the later. In order to substantialize Maldacena's conjecture, a better understanding on the physical contents of the SCFT living on the boundary of AdS space is required. While for the case of AdS_5 , our knowledge is limited on computing the physical correlators in $N = 4$ superconformal Yang-Mills theory, the case of AdS_3 space should provide a better test ground, thanks to the conformal Ward identities of 2d nature, which involves infinitely many conserved charges. Hopefully, the techniques developed in solving the AdS_3/CFT correspondence can be eventually generalized to the cases of higher dimensional AdS spaces.

Presently, there are two main approaches to formulate string theories on AdS_3 space, analogous to the development of the superstring theories by the NSR and GS formalism.

(1) As for the NSR formalism, a quantization procedure on AdS_3 space is given in ref.[4], in which, the spacetime superconformal generators are constructed and some physical states given. The advantage of the NSR formalism is its manifest covariance and worldsheet superconformal invariance. The disadvantage is that the spacetime supersymmetry is not manifest and it is difficult to couple it to the R-R charged background. Another disadvantage is that the physical state content is not manifest and deserves a careful BRST cohomology analysis.

(2) For the case of Green-Schwarz type superstring on AdS_3 space, a κ symmetry invariant action and its simplified (κ symmetry fixed) version are given in refs.[8, 9].³ In the Green-Schwarz formalism, the spacetime supersymmetry is manifest, and both NS-NS and R-R background can be charged. However, due to its high non-linearity, the quantization procedure for the present case is not known, even for its simplified version. Therefore, both worldsheet and spacetime superconformal invariance are not obvious at the quantum level.

The purpose of the present paper is trying to make a connection between the two independent approaches. Our starting point is the NSR formalism. After fixing the conformal gauge, there are still residual conformal gauge symmetries, which are responsible for getting rid of negative norm states and make the string theory unitary. The usual covariant quantization is to enlarge the Hilbert space to include the reparametrization ghosts, and the negative norm states decouple from the physical state space which are BRST invariant. The underlying spacetime field theory should only depend on the physical degrees of freedom, which are transversal to the longitudinal ones. However, there are no prescribed "longitudinal" and "transversal" directions in the Hilbert space, each choice should give the equivalent string theory as ensured by the conformal invariance.

In contrast, in light-cone gauge quantization, the conformal gauge symmetry is completely fixed, and one has to make a definite choice on which are the longitudinal degrees of freedom and which are the transversal ones. In flat spacetime, the light-cone gauge choice is made to

³The nonlinear σ model on coset space $\frac{SU(2,2|4)}{SO(4,1) \times SO(5)}$ are first proposed to describe superstring theory on AdS_5 space. For more details, see refs.[7]. The AdS_3 case is a generalization from that in ref.[7].

align worldsheet “time” direction along a light-cone coordinate in the spacetime. In our AdS_3 case things are more complicated, since, generally in curved spacetime there can be no flat directions along which light moves freely.

In the present paper, we shall propose a way to resolve such kind of problems. The point is that at the AdS_3 boundary, one can find a flat light-cone direction, and at the vicinity of the boundary, the term which curve the space-time is really the screening charge, which can be perturbatively expanded as what are usually done in the calculations of the correlators in WZNW models, cf. ref.[17].

To justify our approach to the quantizations of string theories on AdS_3 space, we have checked the following consistency conditions:

(1) The spacetime superconformal algebra proposed in ref.[4], when re-expressed in our transversal degrees of freedom after the light-cone gauge choice is made, is closed. The conformal anomaly agrees with that given in ref.[4], provided the worldsheet conformal anomalies c equal 26 resp. 15 for bosonic string resp. superstring.

(2) In string theories on AdS_3 space, there is a flat time direction which is dual to the J^3 in $sl(2, R)$ current algebra. Our light-cone gauge choice is the same as choosing this flat time direction combined with another flat space direction to form a light-cone. Since there exist unitary representations in $SL(2, R)/U(1)$ conformal field theory, cf. ref.[12], the unitarity of the string theory is ensured, provided the non- AdS_3 part is made of unitary conformal field theories.

Using the light-cone gauge quantization to get rid of the unphysical degrees of freedom, we recover the spacetime superconformal algebra in terms of the physical transversal degrees of freedom. By this method, we find that the spacetime superconformal field theory is described by the orbifold theory on the target space $\frac{Sym_p(\mathcal{M}_{Liouville} \times S_3 \times T^4)}{\mathbb{Z}^p}$, and spacetime supersymmetry is realized in a nontrivial way. By the similar procedure, the GS superstring theory on $AdS_3 \times S_3 \times T_4$ with NS-NS background [8, 9] can be quantized in the light-cone gauge, giving the same spacetime superconformal algebra as the NSR superstring theory. The generalization of our procedure to the case of R-R charged background is also currently under investigation.[25]

The present paper is organized in the following way. In section 2, we review the NSR formalism of the bosonic string theory on $AdS_3 \times \mathcal{N}$, and give a description of how to make the light-cone gauge choice and what the spacetime conformal field theory is.

In section 3, we generalize the light-cone gauge quantization procedure to the case of fermionic string. A consistent quantization procedure, which makes light-cone gauge choice possible, is given.

In section 4, the explicit construction of the spacetime $N = 4$ $su(2)$ extended superconformal algebra is given in terms of fields living in transverse spacetime.

In section 5, we quantize the Green-Schwarz type superstring theory on $AdS_3 \times S_3 \times T_4$ spacetime with NS-NS background.

In section Conclusions and Discussions, we speculate on how to calculate one loop partition functions and what could be further done using our generalized light-cone gauge quantization procedure.

Appendix A is devoted to clarify our conventions, in particular for the spinors in AdS_3 space, as well as the relation between the $N = 2$ spacetime SUSY algebra $su(1, 1|2)^2$ and the global part of the $N = 4$ spacetime SCA.

2 Bosonic String Theory on AdS_3 Space

2.1 String Action and Light-Cone Gauge Quantization

AdS_3 space is a hypersurface embedded in flat $\mathcal{R}^{2,2}$,

$$-X_{-1}^2 + X_1^2 + X_2^2 - X_3^2 = -l^2 \quad (1)$$

with the induced metric

$$ds^2 = -dX_{-1}^2 + dX_1^2 + dX_2^2 - dX_3^2 \quad (2)$$

It admits the $sl(2, R) \times sl(2, R)$ Lie algebra as its isometry generators,

$$[J_{\alpha\beta}, J_{\gamma\rho}] = \eta_{\alpha\rho} J_{\beta\gamma} + \eta_{\beta\gamma} J_{\alpha\rho} - \eta_{\beta\rho} J_{\alpha\gamma} - \eta_{\alpha\gamma} J_{\beta\rho} \quad (3)$$

where $\eta_{\alpha\beta}$ is the flat metric $diag(-1, 1, 1, -1)$. Let

$$\begin{aligned} J^1 &= -i\frac{1}{2}(J_{23} + J_{-11}) \\ J^2 &= -i\frac{1}{2}(J_{31} + J_{-12}) \\ J^3 &= -i\frac{1}{2}(J_{12} - J_{-13}) \\ \bar{J}^1 &= -i\frac{1}{2}(J_{23} - J_{-11}) \\ \bar{J}^2 &= -i\frac{1}{2}(J_{31} - J_{-12}) \\ \bar{J}^3 &= -i\frac{1}{2}(J_{12} + J_{-13}) \end{aligned} \quad (4)$$

The Commutators eq.(3) become now

$$\begin{aligned} [J^A, J^B] &= i\eta_{CD}\epsilon^{ABC} J^D \\ [\bar{J}^A, \bar{J}^B] &= i\eta_{CD}\epsilon^{ABC} \bar{J}^D \\ [J^A, \bar{J}^B] &= 0 \end{aligned} \quad (5)$$

In conformal field theory, the J^A 's resp. $\bar{J}^A$'s are just the zero modes of the affine $sl(2, R) \times sl(2, R)$ Lie algebra at level k generated by worldsheet holomorphic currents $J^A(z)$ resp. anti-holomorphic ones $\bar{J}^A(\bar{z})$, satisfying the OPE:

$$J^A(z)J^B(w) = \frac{k\eta^{AB}/2}{(z-w)^2} + \frac{i\eta_{CD}\epsilon^{ABC} J^D}{z-w} + \dots \quad (6)$$

where $A, B, C, D=1, 2, 3$ and $\eta_{AB} = diag(+, +, -)$, and similar OPE for the anti-holomorphic ones.

It is convenient to choose the independent coordinates on AdS_3 space as

$$\begin{aligned} \gamma &= \frac{X_2 + iX_1}{X_{-1} + X_3} \\ \bar{\gamma} &= \frac{X_2 - iX_1}{X_{-1} + X_3} \\ \phi &= \frac{\log(X_{-1} + X_3)}{l} \end{aligned} \quad (7)$$

Then the metric tensor becomes

$$ds^2 = l^2(d\phi^2 + e^{2\phi}d\gamma d\bar{\gamma}) \quad (8)$$

As given in ref.[4], the Lagrangian for the bosonic string theory on $AdS_3 \times \mathcal{N}$ with NS-NS background is

$$\mathcal{L} = \partial\phi\bar{\partial}\phi - \frac{2}{\alpha_+}\hat{R}^{(2)}\phi + \beta\bar{\partial}\gamma + \bar{\beta}\partial\bar{\gamma} - \beta\bar{\beta}\exp(-\frac{2}{\alpha_+}\phi) + \mathcal{L}_N \quad (9)$$

where $\alpha_+^2 = 2k - 4$, $k = \frac{l^2}{l_s^2}$, $\mathcal{L}_N$ is the contribution from manifold $\mathcal{N}$. The conformal weights of β, γ are 1, 0 respectively.

The conformal invariant term $\int \beta\bar{\beta}\exp(-\frac{2\phi}{\alpha_+})$ in the action represents the screening charges in the calculations of the correlators in the WZNW theory. Treating it as a perturbation term, then the classical equations of motion for γ, β, ϕ become free ones. The OPE of these fields are

$$\begin{aligned} \phi(z)\phi(0) &= -\log(z) + \dots\dots\dots \\ \beta(z)\gamma(0) &= \frac{1}{z} + \dots\dots\dots \end{aligned} \quad (10)$$

and others are regular.

The $sl(2, R)$ current algebra can be represented by

$$\begin{aligned} J^3 &= \beta\gamma + \frac{\alpha_+}{2}\partial\phi \\ J^+ &= \beta\gamma^2 + \alpha_+\gamma\partial\phi + k\partial\gamma \\ J^- &= \beta \end{aligned} \quad (11)$$

Here, $J^+ = J^1 + iJ^2$, $J^- = J^1 - iJ^2$.

The energy-momentum tensor is given by Sugavara construction:

$$T = -\frac{1}{2}\partial\phi\partial\phi + \beta\partial\gamma - \frac{1}{\alpha_+}\partial^2\phi + T_N \quad (12)$$

where T_N is the energy-momentum tensor coming from the manifold $\mathcal{N}$.

When $\frac{\phi}{\alpha_+}$ is large, we can perturbatively expand the term $e^{\int \beta\bar{\beta}\exp(-\frac{2\phi}{\alpha_+})}$ as screening charges. That means the string is really located in the neighborhood of AdS_3 boundary ($\frac{2\phi}{\alpha_+} \rightarrow \infty$). Module possible contractions with screening charges in the correlators, γ can be treated as a holomorphic field with conformal weight 0. By conformal invariance, we can transform γ into a desired form containing no string excitations. It is convenient to make our light-cone gauge choice as follow:

$$\gamma = e^q z^p \quad (13)$$

Here, the momentum $p = \oint \gamma^{-1}(z)\partial\gamma(z)$ equals the number of times the string winds in the γ space, thus must be an integer. And e^q represents the center of mass coordinate of the string, is not an important factor, since it can always be scaled away by a dilation in z , $z \rightarrow e^{q/p}z$. If the context is clear, we can always set $q = 0$. However, keeping the factor e^q can remind us that γ is in fact a worldsheet scalar, since e^q adds conformal weight p to the vacuum.

Fixing γ we can solve the constraint $T = 0$, where T is defined in eq.(12), for β :

$$\beta(z) = \frac{1}{p}e^{-q}z^{1-p}\left(\frac{1}{2}\partial\phi(z)\partial\phi(z) + \frac{1}{\alpha_+}\partial^2\phi(z) - T_N(z) + \frac{\Delta}{z^2}\right) \quad (14)$$

where Δ is a normal ordering constant.

To justify this as the right light-cone gauge choice and that the transversal degrees of freedom is ghost free, let us first bosonize the β, γ system.

$$\begin{aligned}
\gamma(z) &= e^{\phi_1(z)+\phi_2(z)} \\
\beta(z) &= -\partial\phi_1(z)e^{-\phi_1(z)-\phi_2(z)} \\
\phi(z) &\equiv \phi_3(z) \\
\phi_i(z)\phi_j(w) &= \eta_{ij} \log(z-w) \\
\eta_{ij} &= \text{Diag}\{1, -1, -1\}
\end{aligned} \tag{15}$$

$J^3(z)$ and the stress tensor can be expressed in terms of $\phi_i(z)$ field as

$$J^3(z) = -\partial\phi_2(z) + \frac{1}{2}\alpha_+\partial\phi_3(z) \tag{16}$$

$$T(z) = \frac{1}{2}\eta^{ij}\partial\phi_i(z)\partial\phi_j(z) - \frac{1}{2}\partial^2[\phi_1(z) + \phi_2(z) + \frac{2}{\alpha_+}\phi_3(z)] \tag{17}$$

It is convenient to make an $O(1, 2)$ transformation on the scalar fields,

$$\begin{pmatrix} \Phi_1(z) \\ \Phi_2(z) \\ \Phi_3(z) \end{pmatrix} = \begin{pmatrix} \frac{\sqrt{4+\alpha_+^2}}{2} & \frac{\alpha_+^2}{2\sqrt{4+\alpha_+^2}} & \frac{\alpha_+}{\sqrt{4+\alpha_+^2}} \\ 0 & \frac{2}{\sqrt{4+\alpha_+^2}} & \frac{-\alpha_+}{\sqrt{4+\alpha_+^2}} \\ \frac{1}{2}\alpha_+ & \frac{1}{2}\alpha_+ & 1 \end{pmatrix} \begin{pmatrix} \phi_1(z) \\ \phi_2(z) \\ \phi_3(z) \end{pmatrix} \tag{18}$$

which can be solved to give

$$\begin{pmatrix} \phi_1(z) \\ \phi_2(z) \\ \phi_3(z) \end{pmatrix} = \begin{pmatrix} \frac{\sqrt{4+\alpha_+^2}}{2} & 0 & -\frac{1}{2}\alpha_+ \\ \frac{-\alpha_+^2}{2\sqrt{4+\alpha_+^2}} & \frac{2}{\sqrt{4+\alpha_+^2}} & \frac{1}{2}\alpha_+ \\ \frac{-\alpha_+}{\sqrt{4+\alpha_+^2}} & \frac{-\alpha_+}{\sqrt{4+\alpha_+^2}} & 1 \end{pmatrix} \begin{pmatrix} \Phi_1(z) \\ \Phi_2(z) \\ \Phi_3(z) \end{pmatrix} \tag{19}$$

In terms of $\Phi_i(z)$ fields we have

$$J^3(z) = -\sqrt{\frac{k}{2}}\partial\Phi_2(z) \tag{20}$$

$$T(z) = \frac{1}{2}\eta^{ij}\partial\Phi_i(z)\partial\Phi_j(z) - \frac{1}{\alpha_+}\partial^2\Phi_3(z) + T_N(z) \tag{21}$$

$$\gamma(z) = e^{\sqrt{\frac{2}{k}}(\Phi_1(z)+\Phi_2(z))} \tag{22}$$

$$\beta(z) = (\sqrt{\frac{k}{2}}\partial\Phi_1(z) - \frac{1}{2}\alpha_+\partial\Phi_3(z))e^{\sqrt{\frac{2}{k}}(-\Phi_1(z)-\Phi_2(z))} \tag{23}$$

And the screening charge becomes now.

$$V_+ \equiv \oint dz \beta(z) e^{-\frac{2}{\alpha_+}\phi(z)} = \oint dz \sqrt{\frac{k}{2}}\partial\Phi_1(z) e^{-\sqrt{\frac{2}{\alpha_+}}\Phi_3(z)} \tag{24}$$

where we have dropped a total derivative term.

From the stress tensor eq.(21), we see that there are two flat spacetime directions, Φ_1 and Φ_2 , the former is space-like, and the later, which is dual to the current J^3 , is time-like. Using the residual conformal invariance in the conformal gauge, we can transform the worldsheet “time” direction parallel to the coordinate $\Phi^+(z, \bar{z}) \equiv \Phi_1(z, \bar{z}) + \Phi_2(z, \bar{z})$, i.e.

$$\Phi^+(z, \bar{z}) = 2q^+ + 2p^+\tau \quad (25)$$

From eq.(18) we have $\phi_1 + \phi_2 \propto \Phi_1 + \Phi_2$. We find that making $\Phi_1 + \Phi_2$ the light-cone coordinate is equivalent to making $\gamma(z)$ in the form given by eq.(13), where $\phi^+ = \phi_1 + \phi_2$ is fixed to be $\phi^+ = 2q + 2p\tau$. It is well known that there exist unitary representations in $SL(2, R)/U(1)$ coset space theory with $U(1)$ corresponding to the time-like direction [12]. We conclude that the transversal physical space is ghost free provided the conformal field theory from $\mathcal{N}$ is unitary.

2.2 Spacetime Virasoro Algebra in Light-Cone Gauge

At the AdS_3 boundary, $\phi \rightarrow \infty$, the action given by eq.(9) is invariant under spacetime conformal transformations:

$$\gamma \rightarrow f(\gamma) \quad (26)$$

$$\bar{\gamma} \rightarrow \bar{f}(\bar{\gamma}) \quad (27)$$

$$\beta \rightarrow \beta f'(\gamma)^{-1} + \partial\phi f''(\gamma)^{-1} \quad (28)$$

$$\bar{\beta} \rightarrow \bar{\beta} \bar{f}'(\bar{\gamma})^{-1} + \bar{\partial}\phi \bar{f}''(\bar{\gamma})^{-1} \quad (29)$$

$$\phi \rightarrow \phi - \frac{\alpha_+}{2} \log(f'(\gamma) \bar{f}'(\bar{\gamma})) \quad (30)$$

The Virasoro algebra, generated by the conformal transformations on the boundary of AdS_3 space, is constructed in ref.[4] as

$$L_n = - \oint dz [(1 - n^2) J^3 \gamma^n - \frac{n(n-1)}{2} J^- \gamma^{n+1} + \frac{n(n+1)}{2} J^+ \gamma^{n-1}] \quad (31)$$

The spacetime conformal transformations act on the physical space of the string theory, which are invariant under worldsheet conformal transformations. Thus, it is possible to express the spacetime Virasoro generators in terms of fields in transversal dimensions. Substituting our light-cone gauge choice, eqs.(13, 14) into eq.(31) for the spacetime Virasoro generators, we have

$$\begin{aligned} L_n &= \oint dz \left[\frac{1}{p} \left(-\frac{1}{2} \partial\phi \partial\phi - \frac{1}{\alpha_+} \partial^2\phi + T_N - \frac{\Delta}{z^2} \right) e^{nq} z^{np+1} - \frac{\alpha_+}{2} (n+1) \partial\phi e^{nq} z^{np} \right] \\ &= \oint dz \left\{ \frac{1}{p} \left[-\frac{1}{2} \partial\phi \partial\phi - \left(\frac{1}{\alpha_+} - \frac{\alpha_+}{2} \right) \partial^2\phi + T_N - \frac{\Delta}{z^2} \right] e^{nq} z^{np+1} - \frac{\alpha_+(p-1)}{2p} \partial\phi e^{nq} z^{np} \right\} \end{aligned} \quad (32)$$

Using the OPE of T_N and ϕ , it is ready to check that that the L_n 's in the light-cone gauge constructed above close a Virasoro algebra

$$[L_n, L_m] = n(n^2 - 1) \frac{pc}{12} \delta_{m+n,0} + (n-m) [L_{n+m} + \delta_{n+m,0} \left(\frac{a}{2} + \frac{\Delta}{p} \right)] \quad (33)$$

with

$$c = c_N + 1 + 12\left(\frac{1}{\alpha_+} - \frac{\alpha_+}{2}\right)^2 \quad (34)$$

$$a = \frac{c_N + 1 + \frac{12}{\alpha_+^2}}{12}\left(p - \frac{1}{p}\right) - (p - 1) \quad (35)$$

where c_N is the central charge of T_N .

If we shift L_0 by $\frac{a}{2} + \frac{\Delta}{p}$, eq.(33) becomes the standard form of the Virasoro algebra with central charge cp .

In covariant quantization, the central charge of this spacetime Virasoro algebra is $6kp$, so to get rid of spacetime anomaly we must set $cp = 6kp$. This condition fixes the worldsheet center charge for the string theory to be

$$c_{string} = c_N + 3 + 12\frac{1}{\alpha_+^2} = 26 \quad (36)$$

where $\alpha_+^2 = 2k - 4$.

That means when the string propagates in critical dimension, its light-cone gauge quantization could be equivalent to the covariant one. In critical dimension, the light-cone gauge fixed Virasoro generators should close the same algebra as the covariant one, i.e. the shift of L_0 should be 0, or $\Delta = -\frac{pa}{2}$. Then the Virasoro generators of spacetime CFT can be written as

$$L_n = \frac{1}{p}\tilde{L}_{np} + \frac{c}{24}\left(p - \frac{1}{p}\right)\delta_{n,0} \quad (37)$$

where

$$\tilde{L}_m = \oint dz [T_N - \frac{1}{2}\partial\phi\partial\phi - (\frac{1}{\alpha_+} - \frac{\alpha_+}{2})\partial^2\phi]z^{m+1} \quad (38)$$

We shall show below that eq.(37) defines a conformal transformation: $z \rightarrow w = z^{\frac{1}{p}}$

$$\begin{aligned} T(z) &= (w'(z))^2 T(w) + \frac{c}{12} \left[\frac{w'''(z)}{w'(z)} - \frac{3}{2} \left(\frac{w''(z)}{w'(z)} \right)^2 \right] \\ &= \frac{1}{p^2} z^{\frac{2}{p}-2} T(w) + \frac{c}{24} \left(1 - \frac{1}{p^2} \right) z^{-2} \end{aligned} \quad (39)$$

Only the term with integer power of z are well defined on the z complex plane closing a Virasoro Algebra. Hence

$$\begin{aligned} L_n &= \int dz T(z) z^{n+1} \\ &= \frac{1}{p} \int dw T(w) w^{pn+1} + \frac{c}{24} \left(p - \frac{1}{p} \right) \delta_{n,0} \\ &= \frac{1}{p} \tilde{L}_{pn} + \frac{c}{24} \left(p - \frac{1}{p} \right) \delta_{n,0} \end{aligned} \quad (40)$$

which agrees with eq.(37).

In fact, eq.(37) suggests that the spacetime CFT is an orbifold theory on the target space $\frac{Sym_p \mathcal{M}}{Z^p}$, see subsection 4.2 for detailed discussions.

3 Light-Cone Gauge Quantization of Fermionic String Theory on AdS_3

The light-cone gauge quantization for bosonic string theory on AdS_3 space discussed in the previous section is ready to be generalized to the fermionic case. For fermionic string theory on the group manifold, the worldsheet supersymmetry is achieved by considering super WZNW theory [13]. For each bosonic current, we associate a fermionic partner, which is a worldsheet real fermion field and living in the adjoint representation of the finite dimensional Lie algebra.

For fermionic string theory on AdS_3 space, the affine $sl(2, R)$ currents pertaining to the AdS_3 space can be written as

$$J^A = j^A - i \frac{1}{k} \epsilon_{BC}^A \psi^B \psi^C \quad (41)$$

which satisfy OPE eq.(6). Here we use the convention as in ref. [4]

$$\langle \psi^A(z) \psi^B(w) \rangle = \frac{k \eta^{AB}/2}{z - w} \quad (42)$$

The worldsheet supercurrent G and the energy-momentum tensor are

$$\begin{aligned} G(z) &= \frac{2}{k} (\eta_{AB} \psi^A j^B - \frac{i}{3k} \epsilon_{ABC} \psi^A \psi^B \psi^C) + G_N \\ T(z) &= \frac{1}{k} (j^A j_A - \psi^A \partial \psi_A) + T_N \end{aligned} \quad (43)$$

Set

$$\begin{aligned} J^+ &= J^1 + iJ^2 & J^- &= J^1 - iJ^2 \\ j^+ &= j^1 + ij^2 & j^- &= j^1 - ij^2 \\ \psi^+ &= \psi^1 + i\psi^2 & \psi^- &= \psi^1 - i\psi^2 \end{aligned} \quad (44)$$

j^A 's can be represented as in eq.(11), but now with $\alpha_+^2 = 2k$.

For light-cone gauge quantization, it is convenient to use another set of free fermion fields

$$\begin{aligned} \tilde{\psi}^3 &= \psi^3 - \psi^- \gamma \\ \tilde{\psi}^+ &= \psi^+ + \psi^- \gamma^2 - 2\psi^3 \gamma \\ \tilde{\psi}^- &= \psi^- \end{aligned} \quad (45)$$

And define

$$\begin{aligned} \tilde{\beta} &= \beta - \frac{2}{k} \psi^- \psi^3 \\ &= \beta - \frac{2}{k} \tilde{\psi}^- \tilde{\psi}^3 \end{aligned} \quad (46)$$

We then have

$$\begin{aligned} \langle \tilde{\psi}^3(z) \tilde{\psi}^3(w) \rangle &= \langle \psi^3(z) \psi^3(w) \rangle = \frac{-k/2}{z - w} \\ \langle \tilde{\psi}^+(z) \tilde{\psi}^-(w) \rangle &= \langle \psi^+(z) \psi^-(w) \rangle = \frac{k}{z - w} \end{aligned} \quad (47)$$

and

$$\begin{aligned}\langle \tilde{\beta}(z)\tilde{\psi}^A(w) \rangle &= 0 \\ \langle \tilde{\beta}(z)\gamma(w) \rangle &= \frac{1}{z-w}\end{aligned}\quad (48)$$

Using these fields, the supercurrent and the energy-momentum tensor for the worldsheet CFT can be written as

$$\begin{aligned}G &= \frac{2}{k}(\frac{1}{2}\tilde{\psi}^+\tilde{\beta} - \frac{\alpha_+}{2}\tilde{\psi}^3\partial\phi + \frac{k+2}{2}\tilde{\psi}^-\partial\gamma) + G_N \\ T &= (\tilde{\beta}\partial\gamma - \frac{1}{2}\partial\phi\partial\phi - \frac{1}{\alpha_+}\partial^2\phi) - \frac{1}{k}(-\tilde{\psi}^3\partial\tilde{\psi}^3 + \frac{1}{2}\tilde{\psi}^+\partial\tilde{\psi}^- + \frac{1}{2}\tilde{\psi}^-\partial\tilde{\psi}^+) + T_N\end{aligned}\quad (49)$$

The spacetime Virasoro algebra for fermionic string theory on AdS_3 space is defined in ref.[4]

$$\begin{aligned}L_n &= -\oint dz \oint dw G(z)[(1-n^2)\psi^3\gamma^n + \frac{n(n-1)}{2}\psi^-\gamma^{n+1} + \frac{n(n+1)}{2}\psi^+\gamma^{n-1}]_{(w)} \\ &= -\oint dw[(1-n^2)J^3\gamma^n + \frac{n(n-1)}{2}J^-\gamma^{n+1} + \frac{n(n+1)}{2}J^+\gamma^{n-1}]_{(w)}\end{aligned}\quad (50)$$

Here

$$\begin{aligned}J^3 &= j^3 + \frac{\psi^+\psi^-}{k} = (\tilde{\beta}\gamma + \frac{\alpha_+}{2}\partial\phi) + \frac{\tilde{\psi}^+\tilde{\psi}^-}{k} \\ J^+ &= j^+ + \frac{2\psi^+\psi^3}{k} = (\tilde{\beta}\gamma^2 + \alpha_+\gamma\partial\phi + k\partial\gamma) + \frac{2}{k}\tilde{\psi}^+(\tilde{\psi}^-\gamma + \tilde{\psi}^3) \\ J^- &= j^- - \frac{2\psi^-\psi^3}{k} = \tilde{\beta}\end{aligned}\quad (51)$$

In light-cone gauge quantization, similar to what we have done in the bosonic case, we set

$$\gamma = e^q z^p \quad (52)$$

Making our gauge choice worldsheet supersymmetry invariant requires the OPE between G and γ be regular, so at the same time we set

$$\tilde{\psi}^+ = 0 \quad (53)$$

Now we can solve the constraints $G = 0, T = 0$, where G, T are defined in eq.(44), for $\tilde{\psi}^-$ and $\tilde{\beta}$.

$$\begin{aligned}\tilde{\psi}^- &= \frac{e^{-q}z^{1-p}}{p(k+2)}[-kG_N + \alpha_+\tilde{\psi}^3\partial\phi] \\ \tilde{\beta} &= \frac{e^{-q}z^{1-p}}{p}[\frac{1}{2}\partial\phi\partial\phi + \frac{1}{\alpha_+}\partial^2\phi - \frac{1}{k}\tilde{\psi}^3\partial\tilde{\psi}^3 - T_N + \frac{\Delta}{z^2}]\end{aligned}\quad (54)$$

Substituting eq.(54) into eq.(50), the spacetime Virasoro generators become

$$L_n = \oint dz[\frac{1}{p}(-\frac{1}{2}\partial\phi\partial\phi - \frac{1}{\alpha_+}\partial^2\phi + T_M + \frac{1}{k}\tilde{\psi}^3\partial\tilde{\psi}^3 - \frac{\Delta}{z^2})e^{nq}z^{np+1} - \frac{\alpha_+}{2}(n+1)\partial\phi e^{nq}z^{np}] \quad (55)$$

Comparing eq.(55) for the fermionic string theory with eq.(32) for the bosonic case, we find one more term containing $\tilde{\psi}^3$ coming from AdS_3 part, and $\alpha_+^2 = 2k$ now.

Similar to the bosonic case, if we require the L_n defined in eq.(55) close the same spacetime algebra as the covariant one with central charge $c = 6kp$, then we must have

$$c_{string} = c_N + 3 + 12\left(\frac{1}{\alpha_+^2}\right) + \frac{3}{2} = 15 \quad (56)$$

$$\Delta = -\frac{1}{2}\left[\frac{c_N + 1 + \frac{12}{\alpha_+^2} + \frac{1}{2}}{12}(p^2 - 1) - (p^2 - p)\right] \quad (57)$$

That agrees with the fact that the critical dimension of Fermionic string is 10.

4 $N = 4$ Spacetime SCFT

For the superstring theory on AdS_3 space, because of spacetime supersymmetry, the CFT on the AdS_3 boundary is in fact a superconformal field theory (SCFT). For definiteness, we consider the spacetime manifold to be $AdS_3 \times S^3 \times T^4$ as an example. The spacetime $N = 2$ supersymmetry algebra coming from $AdS_3 \times S^3$ is the global part of the $N = 4$ $su(2)$ extended superconformal algebra [10, 11] for the SCFT on the AdS_3 boundary. In the following subsections, we shall first construct the $N = 4$ SCA explicitly in term of physical transversal degrees of freedom, and then find out the target space in which a possible $N = 4$ nonlinear σ models may live.

4.1 Construction of $N = 4$ Superconformal Currents

The affine $su(2)$ algebra in the spacetime $N = 4$ SCFT is lifted from the worldsheet $su(2)$ currents pertaining to the S_3 group manifold. For $SU(2)$ super WZNW theory[13], the worldsheet current algebra can be represented as

$$K^A = k^A + \frac{1}{2}\psi_\alpha \sigma_{\alpha\beta}^A \bar{\psi}_\beta \quad (58)$$

$$k^A(z)k^B(w) = \frac{(k' - 2)/2 \delta^{AB}}{(z - w)^2} + \frac{i\epsilon^{ABC}k^C}{z - w} + \dots \quad (59)$$

$$K^A(z)K^B(w) = \frac{k'/2 \delta^{AB}}{(z - w)^2} + \frac{i\epsilon^{ABC}K^C}{z - w} + \dots$$

Here, k^A 's are orthogonal to the ψ^A fields. The zero modes of the affine $su(2)$ currents $K^A(z)$'s represent one factor of the $SO(4)$ symmetry on the manifold S^3 . The spacetime affine $su(2)$ algebra is defined in ref.[4]

$$K_n^A = \oint K^A(z) \gamma^n dz \quad (60)$$

In the light-cone gauge, it becomes simply

$$K_n^A = \oint K^A(z) e^{nq} z^{pn} dz = e^{nq} \tilde{K}_{np}^A \quad (61)$$

It is clear that K_n^A 's form a affine $su(2)$ Lie algebra with level $k_{space} = pk'$.

In eqs.(55,61), we have constructed the spacetime Virasoro generators L_n and the affine $su(2)$ Lie algebra in the light-cone gauge. In order the spacetime $N = 4$ SCA close, we still need to construct the spacetime supercurrents in the light-cone gauge. In ref.[4], the global part of the $N = 4$ spacetime superconformal algebra is constructed by covariant quantization, which involves worldsheet superconformal ghosts and picture changing techniques. The full $N = 4$ superconformal algebra can be generated by commuting G_r^α 's with either L_n or the J_n^a 's. However, such construction can not be converted into constructing the supercurrents in the light-cone gauge. The reason is in the later case, the spacetime supercurrents should involve only the physical degrees of freedom, into which we do not know yet how to transform the worldsheet ghost part.

Now it is clear that if our light-cone gauge choice, eqs.(52-54), works, we should be able to construct the spacetime supercurrents in the $N = 4$ SCFT in terms of the transversal degrees of freedom. The superconformal algebra should close with the Virasoro generators and the affine $su(2)$ algebra given by eqs.(55,61). It turns out that indeed there exists a unique expression for the spacetime supercurrents and now we proceed to construct them.

First, we set $q = 0$ and $p = 1$, that means the bosonic part of target space is just one copy of $\mathcal{M}_{Liouville} \times S^3 \times T^4$. The $p \neq 1$ case can be generated from the case $p = 1$ just as in the bosonic string theory.

It is well known that the $N = 4$ SCA admits a $so(4) = su(2) \times su(2)$ automorphism [14, 10]. In our case, the first $su(2)$ comes from the S_3 part, the second $su(2)$ comes from one factor $su(2)$ of the $so(4)$ acting on T^4 . We first decompose our $N = 4$ SCA into representations of the $su(2) \times su(2)$ automorphism. The Virasoro generators, eq.(55) belongs to $(\mathbf{0}, \mathbf{0})$ and the affine $su(2)$ generators, eq.(61) are in the $(\mathbf{1}, \mathbf{0})$. The supercurrent G_α 's belong to the $(\frac{1}{2}, \frac{1}{2})$. To see the action of the $su(2) \times su(2)$ on the G_α 's, it is convenient to write G_α 's in terms of real components $\tilde{G}^A$'s

$$\begin{aligned} G_1 &= \tilde{G}^0 + i\tilde{G}^3 \\ G_2 &= \tilde{G}^2 + i\tilde{G}^1 \end{aligned} \quad (62)$$

Define a matrix (G^{AB}) to be

$$G = i\tilde{G}^A \sigma_A = \begin{pmatrix} \tilde{G}^0 + i\tilde{G}^3 & \tilde{G}^2 + i\tilde{G}^1 \\ \tilde{G}^2 + i\tilde{G}^1 & \tilde{G}^0 - i\tilde{G}^3 \end{pmatrix} = \begin{pmatrix} G_1 & -\bar{G}_2 \\ G_2 & \bar{G}_1 \end{pmatrix} \quad (63)$$

Here, σ^i , $i = 1, 2, 3$ are the Pauli matrices,

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (64)$$

σ^0 is proportional to identity matrix,

$$\sigma^0 = \begin{pmatrix} i & 0 \\ 0 & i \end{pmatrix} \quad (65)$$

and $\sigma_A = (\sigma^A)^*$. Then it is clear that the first $su(2)$ acts on G by left group action, and the second $su(2)$ by right.

Now we also need to decompose the fields living in the transversal space into the representations of the $su(2) \times su(2)$ automorphism. The Liouville field ϕ and its super partner $(\tilde{\psi}^3)$

from the AdS_3 is in the $(\mathbf{0}, \mathbf{0})$. The K^A 's and their super partner χ^A 's are in the $(\mathbf{1}, \mathbf{0})$. Finally, from T^4 , we have the $u(1)$ currents ∂X^N 's and their super partner λ^N 's for $N = 1, 2, 3$ in the $(\mathbf{0}, \mathbf{1})$ and $\partial X^0, \lambda^0$ in the $(\mathbf{0}, \mathbf{0})$. Since the spacetime supercurrents G_α 's are spacetime spinors, and the worldsheet fermions are spacetime vectors, we first need to transform the later into spacetime spinors. This can be done by the usual bosonization procedure.

Define

$$\begin{aligned}
\chi^3 + \sqrt{\frac{2}{k}}\tilde{\psi}^3 &= e^{\phi^1} \\
\chi^3 - \sqrt{\frac{2}{k}}\tilde{\psi}^3 &= e^{-\phi^1} \\
\chi^+ &= e^{\phi^2} \\
\chi^- &= e^{-\phi^2} \\
\lambda^3 + i\lambda^0 &= e^{\phi^3} \\
\lambda^3 - i\lambda^0 &= e^{-\phi^3} \\
\lambda^1 + i\lambda^2 &= e^{\phi^4} \\
\lambda^1 - i\lambda^2 &= e^{-\phi^4}
\end{aligned} \tag{66}$$

Then we can construct the $su(2) \times su(2)$ generators acting on these fermions in term of ϕ^A 's

$$\begin{aligned}
j^+ &= \oint e^{\phi^2}(e^{\phi^1} + e^{-\phi^1}) \\
j^- &= \oint (e^{\phi^1} + e^{-\phi^1})e^{-\phi^2} \\
j^3 &= \oint \partial\phi^2 \\
k^+ &= \oint e^{\phi^4}(e^{\phi^3} + e^{-\phi^3}) \\
k^- &= \oint (e^{\phi^3} + e^{-\phi^3})e^{-\phi^4} \\
k^3 &= \oint \partial\phi^4
\end{aligned} \tag{67}$$

Using the ϕ^A fields and considering the representations of the j^a 's and k^a 's, we can construct a set of worldsheet fermion fields which are also spacetime spinors

$$\begin{aligned}
\psi_1 &= e^{\frac{1}{2}(\phi_1+\phi_2+\phi_3+\phi_4)} \\
\bar{\psi}_1 &= e^{\frac{1}{2}(-\phi_1-\phi_2-\phi_3-\phi_4)} \\
\psi_2 &= e^{\frac{1}{2}(-\phi_1-\phi_2+\phi_3+\phi_4)} \\
\bar{\psi}_2 &= e^{\frac{1}{2}(\phi_1+\phi_2-\phi_3-\phi_4)} \\
\chi_1 &= e^{\frac{1}{2}(-\phi_1+\phi_2+\phi_3-\phi_4)} \\
\bar{\chi}_1 &= e^{\frac{1}{2}(\phi_1-\phi_2-\phi_3+\phi_4)} \\
\chi_2 &= e^{\frac{1}{2}(\phi_1-\phi_2+\phi_3-\phi_4)} \\
\bar{\chi}_2 &= e^{\frac{1}{2}(-\phi_1+\phi_2-\phi_3+\phi_4)}
\end{aligned} \tag{68}$$

The OPE of ψ_α, χ_α are

$$\psi_\alpha(z)\bar{\psi}_\beta(w) = \frac{\delta_{\alpha\beta}}{z-w} + \dots \tag{69}$$

$$\chi_\alpha(z)\bar{\chi}_\beta(w) = \frac{\delta_{\alpha\beta}}{z-w} + \dots \tag{70}$$

and all others are regular.

For free fermions χ_α in $(\frac{1}{2}, \frac{1}{2})$, and $u(1)$ currents in $(\mathbf{0}, \mathbf{0})$ and $(\mathbf{0}, \mathbf{1})$ from T^4 part, it is easy to construct a $N = 4$ SCA with $c = 6$, cf.[10]

$$\begin{aligned}
\hat{G}_1 &= \chi_1 \partial(X^3 + iX^0) - \bar{\chi}_2 \partial(X^1 + iX^2) \\
\bar{\hat{G}}_1 &= \bar{\chi}_1 \partial(X^3 - iX^0) + \chi_2 \partial(X^1 - iX^2) \\
\hat{G}_2 &= \chi_2 \partial(X^3 + iX^0) + \bar{\chi}_1 \partial(X^1 + iX^2) \\
\bar{\hat{G}}_2 &= \bar{\chi}_2 \partial(X^3 - iX^0) + \chi_1 \partial(X^1 - iX^2) \\
\hat{K}^A &= \frac{1}{2} \chi \sigma^A \bar{\chi} \\
\hat{T} &= -\frac{1}{2} \partial X^N \partial X_N + \frac{1}{2} (\partial \chi_1 \bar{\chi}_1 - \chi_1 \partial \bar{\chi}_1 + \partial \chi_2 \bar{\chi}_2 - \chi_2 \partial \bar{\chi}_2)
\end{aligned} \tag{71}$$

Now we are ready to construct another $N = 4$ SCA, which are orthogonal to the one given in eqs.(71), with the bosonic fields coming from $AdS_3 \times S_3$ and free fermions ψ_α 's defined in eq.(68). Let us first consider all the possibilities for constructing G_1 up to a normalization factor

$$G_1 = \psi_1 k^3 + A \psi_2 k^+ - B \partial \psi_1 + C \psi_1 \bar{\psi}_2 \psi_2 + D \psi_1 \partial \phi \tag{72}$$

Here, A,B,C,D are numeric coefficients to be determined.

$K^+(z)G_1(w)$ regular determines $A = 1$. $G_1(z)G_1(w)$ regular determines $C = 1$, $D = (\frac{k'}{2})^{\frac{1}{2}}$. Now consider the OPE $K^-(z)G_1(w)$. By demanding that only simple pole appear determines $B = k' - 1$. Now G_1 is determined up to a normalization factor. By $su(2) \times su(2)$ automorphism, G_2 and $\bar{G}_1, \bar{G}_2$ can also be determined. Finally we have

$$\begin{aligned}
G_1 &= \sqrt{\frac{2}{k'}} [\psi_1 k^3 + \psi_2 k^+ - (k' - 1) \partial \psi_1 + \psi_1 \bar{\psi}_2 \psi_2 + (\frac{k'}{2})^{\frac{1}{2}} \psi_1 \partial \phi] \\
G_2 &= \sqrt{\frac{2}{k'}} [-\psi_2 k^3 + \psi_1 k^- - (k' - 1) \partial \psi_2 + \psi_2 \bar{\psi}_1 \psi_1 + (\frac{k'}{2})^{\frac{1}{2}} \psi_2 \partial \phi] \\
\bar{G}_1 &= \sqrt{\frac{2}{k'}} [\bar{\psi}_1 k^3 + \bar{\psi}_2 k^- + (k' - 1) \partial \bar{\psi}_1 + \bar{\psi}_2 \psi_2 \bar{\psi}_1 - (\frac{k'}{2})^{\frac{1}{2}} \bar{\psi}_1 \partial \phi] \\
\bar{G}_2 &= \sqrt{\frac{2}{k'}} [-\bar{\psi}_2 k^3 + \bar{\psi}_1 k^+ + (k' - 1) \partial \bar{\psi}_2 + \bar{\psi}_1 \psi_1 \bar{\psi}_2 - (\frac{k'}{2})^{\frac{1}{2}} \bar{\psi}_2 \partial \phi]
\end{aligned} \tag{73}$$

It is easy to check

$$\begin{aligned}
K^A(z)G_\alpha(w) &= \frac{\frac{1}{2}G_\beta(\sigma^A)_{\beta\alpha}}{z-w} + \dots\dots\dots \\
K^A(z)\bar{G}_\alpha(w) &= -\frac{\frac{1}{2}(\sigma^A)_{\alpha\beta}\bar{G}_\beta}{z-w} + \dots\dots\dots
\end{aligned} \tag{74}$$

And

$$\begin{aligned}
K^A(z)K^B(w) &= \frac{\frac{1}{2}(k' - 1)}{(z-w)^2} + \frac{i\epsilon^{ABC}K^C}{z-w} + \dots\dots\dots \\
G_\alpha(z)\bar{G}_\beta(w) &= \frac{2T}{z-w} + \frac{2\sigma_A \partial K^A}{z-w} + \frac{4\sigma_B K^B}{(z-w)^2} + \frac{4(k' - 1)}{(z-w)^3} + \dots\dots\dots \\
T(z)G_\alpha(w) &= \frac{G_\alpha}{z-w} + \frac{3/2 G_\alpha}{(z-w)^2} + \dots\dots\dots
\end{aligned} \tag{75}$$

Here

$$T = \frac{1}{k'} k^A k_A + \frac{1}{2} (\partial\psi_1 \bar{\psi}_1 - \psi_1 \partial\bar{\psi}_1 + \partial\psi_2 \bar{\psi}_2 - \psi_2 \partial\bar{\psi}_2) - \frac{1}{2} \partial\phi \partial\phi + \left(\frac{1}{2k'}\right)^{\frac{1}{2}} (k' - 1) \partial^2 \phi \quad (76)$$

The central charge of this spacetime Virasoro algebra is $c = 6(k' - 1)$.

Considering the direct sum of the two orthogonal $N = 4$ SCAs constructed above, we get our final $N = 4$ SCA on the space $AdS_3 \times S_3 \times T^4$ with the central charge $c = 6k'$, the desired result.

$$G_{\alpha,n}^{tot} = \sqrt{\frac{1}{p}} \oint dz (G_\alpha + \hat{G}_\alpha) e^{qn} z^{pn+\frac{1}{2}} \quad (77)$$

$$\bar{G}_{\alpha,n}^{tot} = \sqrt{\frac{1}{p}} \oint dz (\bar{G}_\alpha + \bar{\hat{G}}_\alpha) e^{qn} z^{pn+\frac{1}{2}} \quad (78)$$

$$K_n^{tot,A} = \oint dz (K^A + \hat{K}^A) e^{qn} z^{pn} \quad (79)$$

$$L_n^{tot} = \frac{1}{p} \oint dz [(T + \hat{T}) e^{qn} z^{pn+1}] + \frac{c}{24} (p - \frac{1}{p}) \delta_{n,0} \quad (80)$$

where $c = 6k'$. The L_n should be equivalent to that in eq.(55), then we get $k' = k$

4.2 Orbifold $Sym_p \mathcal{M}/Z_p$ as the Target Space of $N = 4$ SCFT

Superstring theory on AdS_3 space defines a spacetime $N = 4$ superconformal field theory living on the 2d boundary of AdS_3 . Let us denote the target space of the spacetime superconformal field theory for $p \equiv \oint \gamma^{-1} \partial\gamma = 1$ by $\mathcal{M}$. We shall show in this section that for general integer p , the target space is the orbifold $Sym_p \mathcal{M}/Z_p$, where, Sym_p means symmetric product of p objects, and Z_p is the cyclic group of order p . In the following we sketch the proof.

Let us first consider the $N = 4$ theory on the space $Sym_p \mathcal{M}$, which consists of p copies of subspace $\mathcal{M}$, labeled by $\mathcal{M}^i$, $i = 1, 2, \dots, p$. On each $\mathcal{M}^i$, we have an independent $N = 4$ theory corresponding to $p = 1$, which are constructed by the holomorphic fields living in the transversal space in the light-cone gauge. In our case, those fields are ϕ^i , $K^{A,i}$, ψ_α^i , χ_α^i , $\partial X^{N,i}$, which we denote collectively by $\mathcal{J}^{A,i}$. The generators of the $N = 4$ SCA on $\mathcal{M}^i$, such as T^i , G_α^i , $K^{A,i}$ are collectively denoted by $\mathcal{T}^{A,i}$, with the generators of the total $N = 4$ SCA on $Sym_p \mathcal{M}$ given by

$$\mathcal{T}^A = \sum_i \mathcal{T}^{A,i} \quad (81)$$

Now consider the $N = 4$ theory on $Sym_p \mathcal{M}/Z_p$. The Z_p cyclic group is an Abelian group generated by a single element $\hat{\lambda}$, $\hat{\lambda}^p = identity$. $\hat{\lambda}^j$ acts on the $\mathcal{J}^{A,i}$ by shifting the i index by j units: $\mathcal{J}^{A,i} \rightarrow \mathcal{J}^{A,i+j}$. Gauging by the Z_p action means that the $\mathcal{J}^{A,i}$'s are defined up to Z_p actions. That is to say, $\mathcal{J}^{A,i}$'s admit a Z_p holonomy around a non-contractable loop in z complex plane, which represents the AdS_3 boundary. For example, a possible choice of the boundary condition could be

$$\mathcal{J}^{A,i}(ze^{2\pi i}) = \mathcal{J}^{A,i+j}(z) \quad (82)$$

It turns out that any given choice of j in eq.(82) results in an equivalent theory. Thus it suffices to consider the conjugacy class of the Z_p action, which can be represented by a single element

$\hat{\lambda}$ corresponding to $j = 1$ in eq.(82). It is convenient to diagonalize the Z_p action in eq.(82). Define:

$$\lambda = e^{2\pi i/p} \quad (83)$$

$$K^{A,[j]} = \sum_{i=0}^{p-1} \lambda^{-ij} K^{A,i} \quad (84)$$

$$\partial\phi^{[j]} = \sum_{i=0}^{p-1} \lambda^{-ij} \partial\phi^i$$

$$\partial X^{N,[j]} = \sum_{i=0}^{p-1} \lambda^{-ij} \partial X^{N,i}$$

$$\psi_\alpha^{[j]} = \frac{1}{\sqrt{p}} \sum_{i=0}^{p-1} \lambda^{-ij} \psi_\alpha^i$$

$$\chi_\alpha^{[j]} = \frac{1}{\sqrt{p}} \sum_{i=0}^{p-1} \lambda^{-ij} \chi_\alpha^i$$

conversely we have

$$K^{A,i} = \frac{1}{p} \sum_{j=0}^{p-1} \lambda^{ij} K^{A,[j]} \quad (85)$$

$$\partial\phi^j = \frac{1}{p} \sum_{i=0}^{p-1} \lambda^{ij} \partial\phi^{[i]}$$

$$\partial X^{N,j} = \frac{1}{p} \sum_{i=0}^{p-1} \lambda^{ij} \partial X^{N,[i]}$$

$$\psi_\alpha^j = \frac{1}{\sqrt{p}} \sum_{i=0}^{p-1} \lambda^{ij} \psi_\alpha^{[i]}$$

$$\chi_\alpha^j = \frac{1}{\sqrt{p}} \sum_{i=0}^{p-1} \lambda^{ij} \chi_\alpha^{[i]}$$

From eq.(82) by setting $j = 1$, we can read off the mode expansions for $\mathcal{J}^{A,[i]}(z)$

$$\begin{aligned} K^{A,[i]}(z) &= \sum_{n \in \mathbf{Z}} K_{n-i/p}^{A,[i]} z^{-n-1+i/p} \\ \partial\phi^{[i]}(z) &= \sum_{n \in \mathbf{Z}} a_{n-i/p}^{[i]} z^{-n-1+i/p} \\ \partial X^{N,[i]}(z) &= \sum_{n \in \mathbf{Z}} a_{n-i/p}^{N,[i]} z^{-n-1+i/p} \\ \psi_\alpha^{[i]}(z) &= \sum_{n \in \mathbf{Z}} b_{\alpha, n-\frac{1}{2}-i/p}^{[i]} z^{-n+i/p} \\ \chi_\alpha^{[i]}(z) &= \sum_{n \in \mathbf{Z}} c_{\alpha, n-\frac{1}{2}-i/p}^{[i]} z^{-n+i/p} \end{aligned} \quad (86)$$

From the mode expansion in eq.(86), it can be seen that the normalizations in eq.(84) are chosen in such a way that

$$\begin{aligned} [K_{n-i/p}^{A,[i]}, K_{m-j/p}^{B,[j]}] &= f^{ABC} K_{n+m-(i+j)/p}^{C,[i+j]} + \delta_{i+j,0} \delta_{n+m,0} \delta_{a,b} k(pn-i) \\ [b_{\alpha, n+\frac{1}{2}-i/p}^{[i]}, \bar{b}_{\beta, m-\frac{1}{2}-j/p}^{[j]}] &= \delta_{i+j,0} \delta_{n+m,0} \end{aligned} \quad (87)$$

And similar commutation relations for $a_{n-i/p}^{[i]}$, $a_{n-i/p}^{N,[i]}$, $c_{\alpha,n-\frac{1}{2}-i/p}^{[i]}$. Those commutators are to be compared with that of the $N = 4$ theory with the target space $\mathcal{M}$, for which, we have the mode expansions

$$\begin{aligned} K^A(z) &= \sum_{n \in \mathbf{Z}} K_n^A z^{-n-1} \\ \partial\phi(z) &= \sum_{n \in \mathbf{Z}} a_n z^{-n-1} \\ \partial X^N(z) &= \sum_{n \in \mathbf{Z}} a_n^N z^{-n-1} \\ \psi_\alpha(z) &= \sum_{n \in \mathbf{Z}} \psi_{\alpha,n-\frac{1}{2}} z^{-n} \end{aligned} \tag{88}$$

Compare eqs.(88) with eqs.(86), we are led to the following isomorphism:

$$\begin{aligned} K_{n-i/p}^{A,[i]} &\rightarrow K_{pn-i}^A \\ a_{n-i/p}^{[i]} &\rightarrow a_{pn-i} \\ a_{n-i/p}^{N,[i]} &\rightarrow a_{pn-i}^N \\ b_{\alpha,n-\frac{1}{2}-i/p}^{[i]} &\rightarrow b_{\alpha,np-\frac{1}{2}p-i} \\ c_{\alpha,n-\frac{1}{2}-i/p}^{[i]} &\rightarrow c_{\alpha,np-\frac{1}{2}p-i} \end{aligned} \tag{89}$$

Depending on p even or odd, for $N = 4$ theory on $Sym_p \mathcal{M}/Z_p$ with NS boundary condition, we need to define a $N = 4$ theory on $\mathcal{M}$ with Ramand or NS boundary condition.

Now we are ready to express the generators for the total $N = 4$ SCA on $Sym_p \mathcal{M}/Z_p$, eq.(81), in terms of the Z_p twisted fields in eq.(86). Notice that when the fields $\mathcal{J}^{A,[i]}$ are twisted, the zero mode of the energy-momentum tensor acquires a shift, similar to the anomalous transformation of the energy-momentum tensor under the conformal transformation $z \rightarrow z^{\frac{1}{p}}$, cf. eq.(40). For simplicity, let us consider the $T(z)$ in eq.(76) as an example.

$$\begin{aligned} T(z) & \\ = \sum_i T^i + \frac{c}{24}(p - \frac{1}{p})z^{-2} & \\ = \sum_i \frac{1}{k'} k^{A,i} k_A^i + \frac{1}{2}(\partial\psi_\alpha^i \bar{\psi}_\alpha^i - \psi_\alpha^i \partial\bar{\psi}_\alpha^i - \partial\phi^i \partial\phi^i) + (\frac{1}{2k'})^{\frac{1}{2}}(k' - 1)\partial^2\phi^i & \\ = \sum_{i,j,l} \frac{\lambda^{ij}\lambda^{il}}{p^2} (\frac{1}{k'} k^{A,[j]} k_A^{[l]} + \frac{1}{2}(\partial\psi_\alpha^{[j]} \bar{\psi}_\alpha^{[l]} - \psi_\alpha^{[j]} \partial\bar{\psi}_\alpha^{[l]} - \partial\phi^{[j]} \partial\phi^{[l]}) & \\ + \sum_{i,j} \frac{\lambda^{ij}}{p} (\frac{1}{2k'})^{\frac{1}{2}}(k' - 1)\partial^2\phi^{[j]} & \\ = \sum_i \frac{1}{p} \{ \frac{1}{k'} k^{A,[i]} k_A^{[i]} + \frac{1}{2}(\partial\psi_\alpha^{[i]} \bar{\psi}_\alpha^{[i]} - \psi_\alpha^{[i]} \partial\bar{\psi}_\alpha^{[i]} - \frac{1}{2}\partial\phi^{[i]} \partial\phi^{[i]}) \} + (\frac{1}{2k'})^{\frac{1}{2}}(k' - 1)\partial^2\phi^{[0]} & \end{aligned} \tag{90}$$

Substituting eq.(89) into the last line of eq.(90) and making similar changes on $\hat{T}(z)$, we arrive at the eq.(80). Similar relation holds for $G_\alpha(z)$, $K^A(z)$. Thus the $N = 4$ SCA defined in eqs.(77-80) in fact defines a theory on the target space $Sym_p \mathcal{M}/Z_p$.

5 GS Superstring Theory on $AdS_3 \times S_3 \times T_4$ Spacetime with NS-NS Background

In the the section Introduction, we have briefly discussed the difficulty arising, when quantizing the Green-Schwarz type superstring theory on $AdS_3 \times S_3 \times T_4$, due to the high nonlinearity of the equations of motion. In this section, we shall describe how to generalize the light-cone gauge choice to the case of GS string theory on AdS_3 space, in which we get linear equations of motion. In the present paper, we shall only consider the case with pure NSNS background. The RR charged one is more complicated and deserves a separate consideration[25]

The action for the GS string theory on the space $AdS_3 \times S_3 \times T_4$ has been presented in refs.[8, 9, 23]. Here, following ref.[9] and using the convention adopted in Appendix A, we write down the κ symmetry fixed GS superstring action in pure NSNS background.

$$\begin{aligned}
S = & -\frac{1}{2} \int d^2\sigma \{ (\sqrt{g}g^{ij} + \epsilon^{ij})r^2 [\partial_i x^+ + \frac{1}{2}(\bar{\psi}_L \partial_i \psi_L - \partial_i \bar{\psi}_L \psi_L)] \\
& \times [\partial_j x^- + \frac{1}{2}(\bar{\psi}_R \partial_j \psi_R - \partial_j \bar{\psi}_R \psi_R)] \\
& + \sqrt{g}g^{ij} \frac{l^2}{l_s^2} [r^{-2} \partial_i r \partial_j r + tr(\partial_i g \partial_j g^{-1})] + \delta_{rs} \partial_i z^r \partial_j z^s \} \\
& + \frac{i}{3} \int_{M^3} d^3x \frac{l^2}{l_s^2} \epsilon^{ijk} tr(g^{-1} \partial_i g g^{-1} \partial_j g g^{-1} \partial_k g)
\end{aligned} \tag{91}$$

where, z^r are the bosonic fields on T^4 , and g is a group element of $SU(2)$. Using the projective coordinate of S^3 (the group manifold of $SU(2)$), eq.(91) becomes the same action in ref.[9].

In ref.[8], the final form of the action involves only pure RR background and the spinors do not conform to our convention. In the following we shall describe how the κ symmetry is fixed leading to the same action as in eq.(91), in the case when pure NSNS background is considered.

In particular, we shall show here, that the gauge choice in ref.[8]

$$\theta_{R-}^I = \frac{1}{2}(\delta^{IJ} - i\epsilon^{IJ}\gamma_R^0\gamma_R^1)\theta_R^J = 0 \tag{92}$$

is just the light-cone gauge choice for the spacetime spinors. Here the lower index R refers to the gamma matrices and spinors in ref.[8].

Now we rewrite eq.(92) in our convention

$$\begin{aligned}
\theta_-^I &= T^{-1}\theta_{R-}^I \\
&= \frac{1}{2}(\delta^{IJ} - i\epsilon^{IJ}T^{-1}\gamma_R^0\gamma_R^1T)\theta^J \\
&= 0
\end{aligned} \tag{93}$$

i.e.

$$\theta_1^I + i\theta_1^{II} = 0 \quad \theta_2^I - i\theta_2^{II} = 0 \tag{94}$$

And

$$\bar{\theta}_2^I - i\bar{\theta}_2^{II} = 0 \quad \bar{\theta}_1^I + i\bar{\theta}_1^{II} = 0 \tag{95}$$

Set

$$\begin{aligned}\psi_{R\alpha} &= \frac{1}{2}(\theta_\alpha^I - i\theta_\alpha^{II}) & \psi_{L\alpha} &= \frac{1}{2}(\theta_\alpha^I + i\theta_\alpha^{II}) \\ \bar{\psi}_{R\alpha} &= \frac{i}{2}(\bar{\theta}_\alpha^I + i\bar{\theta}_\alpha^{II}) & \bar{\psi}_{L\alpha} &= -\frac{i}{2}(\bar{\theta}_\alpha^I - i\bar{\theta}_\alpha^{II})\end{aligned}\quad (96)$$

Then eqs.(94-95) lead to

$$\Gamma^+ \psi_R = \Gamma^+ \bar{\psi}_L = \Gamma^- \psi_L = \Gamma^- \bar{\psi}_R = 0 \quad (97)$$

where $\Gamma^+ = \Gamma^1 + \Gamma^2$ and $\Gamma^- = \Gamma^1 - \Gamma^2$.

Here we ignore the indices of the spinor representation of $su(2)$. We realize that eq.(97) is the same as fixing the κ symmetry in the light-cone gauge for the spacetime spinors in the flat spacetime, cf. [15].

Using $\psi_R, \psi_L, \bar{\psi}_R, \bar{\psi}_L$ to denote the nonzero component of the spinors, then the action in ref.[8] with pure NSNS background can be written in the same form as given in ref.[9].

To compare eq.(91) with the NSR string, we introduce an auxiliary field β as the Lagrange multiplier. The action, eq.(91), then becomes

$$\begin{aligned}S = & \frac{1}{2} \int dz d\bar{z} \{ -\beta \bar{\beta} \exp(-\frac{2}{\alpha_+} \phi) + \beta [\bar{\partial} x^+ + \frac{1}{2}(\bar{\psi}_L \bar{\partial} \psi_L - \bar{\partial} \bar{\psi}_L \psi_L)] \\ & + \bar{\beta} [\partial x^- + \frac{1}{2}(\bar{\psi}_R \partial \psi_R - \partial \bar{\psi}_R \psi_R)] + \partial \phi \bar{\partial} \phi - \frac{2}{\alpha_+} \hat{R}^{(2)} \phi \} \\ & - \frac{1}{2} \int d^2 \sigma \sqrt{g} g^{ij} [\frac{l^2}{l_s^2} \text{tr}(\partial_i g \partial_j g^{-1}) + \partial_i z^r \partial_j z^r] \\ & + \frac{i}{3} \int_{M^3} d^3 x \frac{l^2}{l_s^2} \epsilon^{ijk} \text{tr}(g^{-1} \partial_i g g^{-1} \partial_j g g^{-1} \partial_k g)\end{aligned}\quad (98)$$

where $\phi = \alpha_+ \log r$ and $\alpha_+^2 = 2k = 2\frac{l^2}{l_s^2}$.

The S^3 part of this action is just a WZNW model in $SU(2)$ group manifold. Because the radiuses of AdS_3 and S^3 are the same, it is obvious that the $sl(2, R)$ and $su(2)$ currents have the same level, as expected from subsection 4.1.

Regarding $\beta \bar{\beta} \exp(\frac{-2\phi}{\alpha_+})$ as a screen charge, we get the the following linearized equations of motion for free bosons and fermions.

$$\begin{aligned}\bar{\partial} \beta &= 0 & \partial \bar{\beta} &= 0 \\ \bar{\partial} \psi_L &= 0 & \partial \psi_R &= 0 \\ \bar{\partial} \bar{\psi}_L &= 0 & \partial \bar{\psi}_R &= 0 \\ \bar{\partial} x^+ &= 0 & \partial x^- &= 0\end{aligned}\quad (99)$$

It is clear now that the light-cone gauge choice for the spinors in the GS type action is self consistent, and we can quantize the action by introducing OPE of free fields. Fixing the κ symmetry is equivalent to fixing the redundant fermionic degrees of freedom, just as what we have done in the light-cone gauge quantization of the NSR string. Now the bosonic part of this action is the same as NSR formulation. So, in $AdS_3 \times S^3 \times T^4$ spacetime, GS string in pure NSNS field background is equivalent to NSR superstring, as can be seen by taking the light-cone gauge. The rest part of quantization procedure goes exactly the same way as the light-cone gauge quantization in the NSR formalism.

6 Conclusions and Discussions

In conclusion, we have presented the light-cone gauge quantization procedure for strings propagating on AdS_3 space charged with NS-NS fields. The procedure works in that we get the right spacetime conformal anomalies for critical bosonic string as well as superstrings. The $N = 4$ spacetime superconformal algebra arises in a natural way for light-cone gauge fixed superstrings, both in NSR and GS formalism. Further, we have proven that in our light-cone gauge choice, the resulting theory is unitary, hence a well define quantum theory.

To quantize superstring theory charged with R-R fields, it seems more appropriate to formulate the string theory in a way with manifest spacetime supersymmetries, as what has been done in refs.[8, 9]. However, the classical equations of motion derived this way are highly nonlinear, posing difficulties in choosing a convenient quantization procedure. The same non-linearity problem in the case of flat spacetime is solved by choosing a light-cone gauge. We believe that in the case of conformal invariant background such as $AdS_3 \times S_3 \times T^4$ or $AdS_5 \times S_5$, a generalized light-cone gauge choice can always be found. The case with NS-NS background worked out in this paper provide just an example, and such quantization procedure should be generalizable to the cases involving R-R charges. Currently, we are making progress along such directions.[25]

In the present paper, we are assuming that the worldsheet theory corresponding to the AdS_3 space is a standard WZNW theory. However for $p > 0$, the holomorphic conformal scalar field γ vanishes at the worldsheet origin, and β , which has conformal weight one, is singular at the origin. In fact, $\gamma(z) \sim z^p$, $\beta(z) \sim z^{-p-1}$ when $z \rightarrow 0$. This is equivalent to the shifting of the Bose sea level, and hence shifting the vacuum state in the WZNW theory, cf. [16]. That would has some effects when coming to the calculations of the correlation functions on the worldsheet. Such point is not elaborated in our present paper and will be left for future discussions.

Note Added: After the completion of the present work, we became aware of the paper, ref.[24], the authors of which found essentially the same target space for the $N = 4$ SCFT as what we presented in Sec.4.2 from a different approach.

Acknowledgment

It is pleasant to acknowledge our discussions with M. Li, Y. Gao and C. Zhu.

This work is supported in part by grants from the NSFC.

Appendix A

The $N = 2$ SUSY algebra $su(1, 1|2)^2$, cf. ref.[8], can be presented as

$$\begin{aligned}
 \{Q_I, \bar{Q}_J\} &= 2\delta_{IJ}(iP_a\gamma^A - P_{a'}\gamma^{a'}) + \epsilon_{IJ}(J_{ab}\gamma^{ab} - J_{a'b'}\gamma^{a'b'}) \\
 [P_a, Q_I] &= -\frac{i}{2}\epsilon_{IJ}\gamma_a Q_J & [P_{a'}, Q_I] &= \frac{1}{2}\epsilon_{IJ}\gamma_{a'} Q_J \\
 [J_{ab}, Q_I] &= -\frac{1}{2}\gamma_{ab} Q_I & [J_{a'b'}, Q_I] &= -\frac{1}{2}\gamma_{a'b'} Q_I \\
 [P_a, \bar{Q}_I] &= \frac{i}{2}\bar{Q}_J\epsilon_{JI}\gamma_a & [P_{a'}, \bar{Q}_I] &= -\frac{1}{2}\bar{Q}_J\epsilon_{JI}\gamma_{a'}
 \end{aligned} \tag{100}$$

$$\begin{aligned}
[J_{ab}, \bar{Q}_I] &= \frac{1}{2} \bar{Q}_I \gamma_{ab} & [J_{a'b'}, \bar{Q}_I] &= \frac{1}{2} \bar{Q}_I \gamma_{a'b'} \\
[J_{AB}, J_{CD}] &= \eta_{BC} J_{AD} + \eta_{AD} J_{BC} - \eta_{AC} J_{BD} - \eta_{BD} J_{AC} \\
[J_{A'B'}, J_{C'D'}] &= \delta_{B'C'} J_{A'D'} + \delta_{A'D'} J_{B'C'} - \delta_{A'C'} J_{B'D'} - \delta_{B'D'} J_{A'C'} \\
P_a &= J_{0a}, & P_{a'} &= J_{0a'}
\end{aligned}$$

where, $Q_{I\alpha\alpha'}$ are the 2 complex chiral 6D spinors, with $I = I, II$, $\alpha = 1, 2$, $\alpha' = 1, 2$. The convention we use here is

$$\begin{aligned}
A, B, C, D &= -1, 1, 2, 3 & \eta_{AB} &= (-1, -1, 1, 1) \\
A', B', C', D' &= 0, 4, 5, 6 & \eta_{A'B'} &= (1, 1, 1, 1) \\
a, b, \dots &= 1, 2, 3 & \eta_{ab} &= (-1, 1, 1) \\
a', b', \dots &= 4, 5, 6 & \eta_{a'b'} &= (1, 1, 1) \\
\gamma^3 &= -\sigma^3 & \gamma^1 &= i\sigma^2 & \gamma^2 &= \sigma^1 \\
\gamma^4 &= \sigma^1 & \gamma^5 &= \sigma^2 & \gamma^6 &= \sigma^3
\end{aligned} \tag{101}$$

In the following we use γ^a , $\gamma^{a'}$ to represent $\gamma^a \otimes 1_2$, $1_2 \otimes \gamma^{a'}$ resp.

The Dirac matrices γ^a used here is a little different from that in ref.[8]. That means we use another basis of the spinors. If we use Q_R representing the SUSY charge in [8], the relation between these two conventions are

$$\begin{aligned}
Q &= T Q_R & \theta &= T^{-1} \theta_R \\
\gamma^3 &= -T \gamma_R^1 T^{-1} & \gamma^1 &= T \gamma_R^3 T^{-1} & \gamma^2 &= T \gamma_R^2 T^{-1}
\end{aligned} \tag{102}$$

where, T is a unitary transformation of the spinors.

$$T = \sqrt{\frac{1}{2}} \begin{pmatrix} -i & -i \\ 1 & -1 \end{pmatrix} \tag{103}$$

From now on we separate the bosonic part algebra $so(2, 2) \times so(4)$ as $(so(1, 2) \times so(3))_L \times (so(1, 2) \times so(3))_R$, and just concentrate ourselves on the left-moving part only.

The algebra can be written as

$$\begin{aligned}
J^1 &= -\frac{i}{2}(J_{23} - J_{-11}) & K^4 &= -\frac{i}{2}(J_{56} + J_{04}) \\
J^2 &= -\frac{i}{2}(J_{31} + J_{-12}) & K^5 &= -\frac{i}{2}(J_{64} + J_{05}) \\
J^3 &= -\frac{i}{2}(J_{12} + J_{-13}) & K^6 &= -\frac{i}{2}(J_{45} + J_{06})
\end{aligned} \tag{104}$$

$$\tag{105}$$

Actually, $so(1, 2)$ is equivalent to $sl(2, R)$, so we can write the algebra as L_0, L_1, L_{-1} .

$$L_0 = iJ^3 \quad L_{\pm 1} = i(J^1 \pm J^2) \tag{106}$$

The super charges can also be combined to form a left-moving one,

$$\begin{aligned}
Q_{\alpha\alpha'} &= \frac{1}{2}(Q_{I\alpha\alpha'} + iQ_{II\alpha\alpha'}) \\
\bar{Q}_{\alpha\alpha'} &= \frac{1}{2}(i\bar{Q}_{I\alpha\alpha'} + \bar{Q}_{II\alpha\alpha'})
\end{aligned} \tag{107}$$

It is easy to check

$$\begin{aligned} [J^a, Q_{\alpha\alpha'}] &= \frac{i}{2}\gamma^a Q_{\alpha\alpha'} & [J^a, \bar{Q}_{\alpha\alpha'}] &= -\frac{i}{2}\bar{Q}_{\alpha\alpha'}\gamma^a \\ [K^{a'}, Q_{\alpha\alpha'}] &= -\frac{1}{2}\gamma^a Q_{\alpha\alpha'} & [K^{a'}, \bar{Q}] &= \frac{1}{2}\bar{Q}_{\alpha\alpha'}\gamma^a \end{aligned} \quad (108)$$

Set

$$\begin{aligned} G_{\frac{1}{2}\alpha'} &= \bar{Q}_{1\alpha'} & \bar{G}_{-\frac{1}{2}\alpha'} &= Q_{1\alpha'} \\ G_{-\frac{1}{2}\alpha'} &= \bar{Q}_{2\alpha'} & \bar{G}_{\frac{1}{2}\alpha'} &= -Q_{2\alpha'} \end{aligned} \quad (109)$$

$L_0, L_1, L_{-1}, K^{a'}$ and $G, \bar{G}$ close an algebra

$$\begin{aligned} [L_0, L_{\pm 1}] &= \mp L_{\pm 1} & [L_{+1}, L_{-1}] &= 2L_0 \\ [L_n, G_r] &= \left(\frac{1}{2}n - r\right)G_{n+m} \\ [G_r, \bar{G}_s] &= 2L_{r+s} + 2(r-s)\sigma_{a'}J_{r+s}^{a'} \end{aligned} \quad (110)$$

which is exactly the left-moving part of the global $N = 4$ SCA in NS sector.

Extending this 6D super algebra to 10D is straightforward. The $(9+1)$ D gamma matrices can be expressed using $\gamma^a, \gamma^{a'}, \gamma^r$.

$$\begin{aligned} \Gamma^a &= \gamma^a \otimes (1_2 \otimes 1_4) \otimes \sigma^1 \\ \Gamma^{a'} &= 1_2 \otimes (\gamma^{a'} \otimes \bar{\gamma}^5) \otimes \sigma^2 \\ \Gamma^r &= 1_2 \otimes (1_2 \otimes \gamma^r) \otimes \sigma^2 \end{aligned} \quad (111)$$

where γ^r ($r=7,8,9,10$) are the 4D gamma matrices, and $\bar{\gamma}^5 = \gamma^7\gamma^8\gamma^9\gamma^{10}$. And the preceding discussions on the relation between the superalgebra $su(1,1|2)^2$ and the global part of the $N = 4$ SCA go over to the 10D case.

References

- [1] J. Maldacena, *The Large N Limit of Superconformal Field Theories and Supergravity*, hep-th/9711200 (1997), Adv.Theor.Math.Phys.2(1998) 231.
- [2] E.Witten, *Anti-de Sitter Space and Holography*, hep-th/9802150 (1998), Adv. Theor. Math. Phys.2(1998), 253.
- [3] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge Theory Correlators from Non-Critical String Theory*, hep-th/9802109 (1998), Phys.Lett. B428 (1998), 105.
- [4] Amit Giveon, David Kutasov and Nathan Seiberg, *Comments on String Theory on AdS_3* , hep-th/9806194 (1998).
- [5] A. Strominger, *Black Hole Entropy from Near-Horizon Microstates*, J. High Energy Phys. 02 (1998) 9, hep-th/9712251
- [6] J. Maldacena and A. Strominger, *AdS_3 Black Holes and a String Exclusion Principle*, hep-th/9804085

- [7] R.R. Metsaev and A.A. Tseytlin, *Type IIB Superstring Action in $AdS_5 \times S^5$ background*, hep-th/9805028 (1998), Nucl.Phys. B533(1998) 109.
R.Kallosh, A.A. Tseytlin, *Simplifying Superstring Action on $AdS_5 \times S^5$* , hep-th/9808088 J.High Energy Phys. 9810(1998) 016.
- [8] J.Rahmfeld and A.Rajaraman, *The GS String Action on $AdS_3 \times S^3$ with Ramond-Ramond Charge*, hep-th/9809164 (1998).
- [9] I.Pesando, *The GS Type IIB Superstring Action on $AdS_3 \times S^3 \times T^4$* , hep-th/9809154 (1998).
- [10] M.Yu, *The Unitary Constructions of the $N=4$ $SU(2)$ Extended Superconformal Algebra*, Phys. Lett. B196(1987) 345.
M.Yu, *The Unitary Representations of the $N=4$ $SU(2)$ Extended Superconformal Algebras*, Nucl. Phys. B294(1987) 890.
- [11] M. Ademollo et al, *Dual String Models with Non-Abelian Color and Flavour Symmetris*, Nucl. Phys. B114 (1976) 297
- [12] L.J. Dixon and M.E. Peskin, *$N=2$ Superconformal Symmetry and $SO(2,1)$ Current Algebra*, Nucl. Phys. B325(1989) 329.
- [13] P. De Vecchia, V.G. Knizhnik, J.L. Petersen and P. Rossi, *A Supersymmetric Wess-Zumino Lagrangian in Two Dimensions*, Nucl. Phys. B253 (1985) 701
- [14] A. Schwimmer, N. Seiberg, *Comments on the $N=2$, $N=3$, $N=4$ Superconformal Algebras in Two-Dimensions*, Phys. Lett. 184B, (1987) 191
- [15] M. Green, J. Schwarz and E. Witten, *Superstring Theory*, Vol. 1, section 5.2.1, Cambridge Univ. Press (1987)
- [16] D. Friedan, E. Martinec and S. Shenker, *Conformal Invariance, Supersymmetry, and String Theory*, Nucl. Phys. B271 (1986) 93
- [17] J. L. Petersen, J. Rasmussen and M. Yu, *Free Field Realizations of 2D Current Algebras, Screening Currents and Primary Fields*, hep-th/9704052 (1997), Nucl.Phys. B502 (1997) 649
- [18] M.J. Duff, H. Lu and C.N. Pope, *$AdS_3 \times S^3$ (Un)twisted and Squashed, and An $O(2,2;\mathbb{Z})$ Multiplet of Dyonic Strings*, hep-th/9807173(1998).
- [19] K. Ito, *Extended Superconformal Algebras on AdS_3* , hep-th/9811002(1998).
- [20] D.Kutasov, F.Larsen and R.G. Leigh, *String Theory in Magnetic Monopole Backgrounds*, hep-th/9812027 (1998).
- [21] J. de Boer, H. Ooguri, H. Robin and J. Tannenhauser, *String Theory on AdS_3* , hep-th/9812046(1998).
- [22] S. Elitzur, O. Feinerman, A. Giveon and D. Tsabar, *String Theory on $AdS_3 \times S^3 \times S^3 \times S_1$* , hep-th/9811245(1998).
- [23] J. Park and S. Rey, *Green-Schwarz Superstring on $AdS_3 \times S^3$* , hep-th/9812062(1998).

- [24] K. Hosomichi and Y. Sugawara, *Hilbert Space of Space-time SCFT in AdS_3 Superstring and T^{4kp}/S_{kp} SCFT*, hep-th/9812100(1998).
- [25] J.Jing, M.Yu, B.Zhang and S.H. Zhao, *Quantization of RR Charged Superstring Theory on AdS_3 Space*, work in progress.